

Research Article | Annual Meeting of the Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA), Miami, Florida, USA, July 14–16, 2016

Practical Assessment of Psoriasis Clinical Severity in both Clinical Trials and Clinical Practice Settings: A Report from the GRAPPA 2016 Annual Meeting

Joseph F. Merola and Alice B. Gottlieb

The Journal of Rheumatology May 2017, 44 (5) 691-692; DOI: <https://doi.org/10.3899/jrheum.170147>

[Article](#)[References](#)[Info & Metrics](#)[PDF](#)[eLetters](#)[← Previous](#)[Next →](#)

Abstract

At the 2016 annual meeting of the Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA), we presented the case for quantitatively assessing the extent of both psoriasis and psoriatic arthritis in the clinical setting, with a particular focus on the validation and expanded novel use of the PGxBSA (static physician's global assessment × body surface area of involvement) in the era of targeted metrics. Herein, we summarize our presentation.

Key Indexing Terms:

[PSORIATIC ARTHRITIS](#)[PSORIASIS](#)[COMBINED CLINIC](#)[GRAPPA](#)[View Full Text](#)

Log in using your username and password